

Nazca plate buoyancy and mantle convection under the Pantanal Basin

J. Assunção¹, V. Sacek¹

¹Instituto de Astronomia, Geofísica e Ciências Atmosféricas da Universidade de São Paulo

The Pantanal wetland is a Quaternary basin in Southwestern Brazil, reaching up to ~500 m of sedimentary package. Although the relatively recent formation of the basin, the mechanisms involved in the origin of the regional subsidence are not fully understood. Previous works proposed a flexural origin for the basin, invoking the last pulse of Andean tectonism and consequent topographic load as the source for the creation of the Pantanal Basin. However, taking into account the flexural rigidity of the South American lithosphere and the Andean geometry, it is difficult to explain the subsidence amplitude as well as the position of the basin due to flexural effects. Therefore, probably another mechanism must be involved in the formation and evolution of this active sedimentary basin. In the present work we will perform several 2D numerical simulations to evaluate how initial conditions can contribute to the buoyancy of the subducting Nazca plate. The results will be used in an upcoming study to calculate the influence of mantle convection generated by the subducting slab geometry and assess if it can induce dynamic topography and contribute to create the depression associated with the Pantanal Basin. We used a finite element code to simulate the thermochemical convection in the mantle, calculating the associated dip angle of subduction along a profile from 80.0°W to 48.5°W, crossing the Pantanal wetland at a latitude of 18°S. To simulate the mantle flow, the thickness of both lithosphere and crust were resampled from the global LITHO1.0 model and the initial thermal structure for subducting slab was constructed based on a simplified thermokinematic model. The subducting slab geometry adopted was derived from the Slab2 model. The results will be considered to three dimensional models in the future to better represent latitudinal variations of the subducting slab and consequently improve the representation of the dynamic topography in the interior of the South American Plate.